

	Vaccine Type	Adult Vaccine Recommendation	Dose(s)
<input type="checkbox"/>	Diphtheria provided as Td or Tdap	<ul style="list-style-type: none"> • Everyone 	Booster every 10 years
<input type="checkbox"/>	Dukoral®	<ul style="list-style-type: none"> • Travellers, depending on destination 	Taken orally, 2 doses taken 1-6 wks apart; last dose taken at least 1 wk before travel Revaccination: <ul style="list-style-type: none"> • If up to 5 years since last dose: 1 dose • If more than 5 years since last dose: 2 doses
<input type="checkbox"/>	Hepatitis A Avaxim®, Havrix®	<ul style="list-style-type: none"> • People with certain medical, occupational, or lifestyle risks • Travellers to high risk destinations 	<ul style="list-style-type: none"> • Avaxim®: initial dose plus 1 follow-up dose 6-36 months later • Havrix®: initial dose plus 1 follow-up dose 6-12 months later
<input type="checkbox"/>	Hepatitis B: Engerix-B® Recombivax HB® Twinrix®	<ul style="list-style-type: none"> • People with certain medical, occupational, or lifestyle risks • Adults living communities or traveling in destinations that are high risk 	<ul style="list-style-type: none"> • 3 doses: initial dose plus 2 follow-up doses at 1 and 6 months later OR • 4 doses (rapid schedule): initial dose plus 3 follow-up doses at 7 days, 21 days & 1 year later OR • 4 doses (accelerated schedule): initial dose plus 3 follow-up doses 1, 2 & 12 months later
<input type="checkbox"/>	Herpes zoster (shingles) non-live vaccine (Shingrix)	<ul style="list-style-type: none"> • Adults 50 years or over; may be given to those 18 years or older at increased risk of herpes zoster due to immunodeficiency or immunosuppression 	<ul style="list-style-type: none"> • Shingrix: 2 doses (2-6 months apart)
<input type="checkbox"/>	Human papillomavirus (HPV) Gardasil®9	<ul style="list-style-type: none"> • Women 18 to 45 years of age • Men 18 to 26 years of age with lifestyle or other risks • Men 27 years of age and older at continued risk of exposure 	3 doses (initial dose plus 2 follow-up doses 2 months and 6 months later)
<input type="checkbox"/>	Influenza (flu)	<ul style="list-style-type: none"> • Adults and children 6 months of age and over 	Annually
<input type="checkbox"/>	Measles/mumps/rubella (German measles) - MMR MMR®II, Priorix®	<ul style="list-style-type: none"> • Adults born before 1970: 1 dose if travelling, have certain medical conditions, or have occupational risk • Adults born in or after 1970 who have not had the disease: 2 doses 	or 2 doses, 4 weeks apart
<input type="checkbox"/>	Meningococcal infection ACYW-135 (Menactra®)	<ul style="list-style-type: none"> • People with specific medical conditions, travellers to certain destinations and those living in communal residences, including students and military personnel 	<ul style="list-style-type: none"> • 1 dose • Revaccination every 5 years if at continued risk
<input type="checkbox"/>	Meningococcal B (Trumenba®)	<ul style="list-style-type: none"> • Adults who are medically high-risk • Adults with occupational risk • Travellers to certain high risk destinations 	<ul style="list-style-type: none"> • An initial dose plus a follow-up dose 6 months later; people at high risk: initial dose plus 2 follow-up doses 1 month and 4-6 months later
<input type="checkbox"/>	Pertussis (whooping cough) provided as Tdap	<ul style="list-style-type: none"> • During each pregnancy 	Once if no prior pertussis vaccination in adulthood
<input type="checkbox"/>	Pneumococcal infection Prevnar®20 Pneumovax®23	<ul style="list-style-type: none"> • Adults 18 yrs and over at risk of pneumonia due to underlying condition • Immunocompromised adults • Adults 65 years of age and over • Adults living in residential care or assisted living facilities • Adults with medical or lifestyle risks 	Prevnar®20: 1 dose Pneumovax®23: 1 dose plus a booster shot in 5 years for those who are immunocompromised
<input type="checkbox"/>	Tetanus (lockjaw) provided as Td or Tdap	<ul style="list-style-type: none"> • Everyone 	Booster every 10 years
<input type="checkbox"/>	Varicella (chickenpox)	<ul style="list-style-type: none"> • People who have not had the disease or been vaccinated 	2 doses, at 4-8 weeks apart depending on vaccine brand
<input type="checkbox"/>	Additional travel vaccines	<ul style="list-style-type: none"> • Vary by destination; consult a travel health clinic or pharmacy with travel health designation, your doctor, or local public health unit 	Vary depending on vaccine
<input type="checkbox"/>	Respiratory syncytial virus or RSV	<ul style="list-style-type: none"> • 60+ or those who are in their 3rd trimester of pregnancy, Wk 32-36 	<ul style="list-style-type: none"> • 1 dose